Table of Contents

1. Executive Summary

Research context

The New Zealand Conservation Trust (NZCT) has recently established a predator trapping programme at Styx Mill Reserve to help contribute to the goal of Predator Free 2050 in New Zealand. The NZCT requires a way to measure conservation progress against.

Aim

To identify the most appropriate ecological biodiversity benchmark to measure bird conservation efforts at Styx Mill Conservation Reserve.

Methodology

- Qualita0 1 227.6 BDC q0.00000887 0 595.2

3.3 Ecological Diversity Benchmarks

There is a vast range of ecological diversity indices, which academics agree can be confusing and often inaccurate (Daly et al., 2018; Hugget, 2005; Redžić, 2007; Reyers et al., 2010; Yen et al., 2019). The most common way to measure biodiversity and develop ecological benchmarks

4. Methods

4.1 Qualitative

A large proportion of the information collected for this report was obtained from the literature reviews. Literature reviews were used to synthesise existing information surrounding the research question, and to understand where further research was required.

Additionally, information was collected by interviewing relevant experts. This included Daniel Leadbeater and Jan Hellyer from NZCT, Julia Rambacher from SLLT and Prof. Dave Kelly and Prof. Jim Briskie from the University of Canterbury. A semi-structured interview style was primarily used, which involved using a list of predetermined questions and allowing open responses from participants.

SMCR was visited to gain an insight into how bird counts are undertaken. During site visits, the bird-count transect was walked. This was to gain a real-life understanding on how data is collected to further improve data collection methods.

4.2 Quantitative methodology

Statistical data analysis and species diversity calculations were used to develop baselines for bird diversity at SMCR. Microsoft Excel was used to analyze trends in the Wetland bird abundance data (Crossland, 2013) and bush bird abundance data (SLLT, 2020). datasets through graphing and modelling techniques.

are relatively stable throughout the years, until a rapid spike in introduced diversity in 2020; this observation corresponds with the increased abundances in introduced species in this time.

It could be extrapolated that the recent establishment of the trapping regime may have led this rise in introduced species, as this was a potential outcome noted regularly throughout the literature (Starling-Windhof et al., 2011). However, without a continuous dataset, other influences cannot be ruled out.

Counts and identification of predators caught since the implementation of the trapping project are recorded in Fig. 3. The results demonstrate the various predatory mammals present at SMCR. The main predator successfully being captured are rats, suggesting they are the most abundant. It is important to note the significant rise in rats in May could be due to the emptying of traps after the COVID-19 lockdown. The recent small peak in possums and decrease in rats recorded in September may suggest a decrease in rat abundances, potentially allowing for other species to interact with traps. This may be a seasonal trend, which could be determined by future years data.

Fig. 3. Trapping data for the recently implemented regime at SMCR. Adapted from NZCT trap emptying data. Updated 6th September by Jan Hellyer.

6. Discussion

6.1 Indicator Species

The initial goal for the trapping project, as communicated by the NZCT, was for the trapping to increase the abundance of all bird species at SMCR. However, through discussions with experts and a review of the current literature, this goal was considered too general and undefined. Although increasing overall abundance at the SMCR may be positive, an increase could be due

Fig. 8. Map of the Styx Mill Reserve with the current line transect as well as the additional proposed line transect and point count locations. Figure generated via Google Earth by Cummack, E.

7.5 Community engagement

Kaval, P., & Roskruge, M. J. (2009).

Ruffell, J., & Didham, R. (2016). Conserving biodiversity in New Zealand's lowland landscapes: Does forest cover or pest control have a greater effect on native birds? *New Zealand Journal of Ecology, 41*(1). doi:10.20417/nzjecol.41.12

Russell, J. (2014). A comparison of attitudes towards introduced wildlife in New Zealand in 1994 and 2012. *Journal of the Royal Society of New Zealand*, *44*(4), 136-151. doi: 10.1080/03036758.2014.944192

Russell, J., Innes, J., Brown, P., & Byrom, A. (2015). Predator-Free New Zealand: Conservation Country. *BioScience,65*(5), 520-525. Retrieved August 10, 2020, from www.jstor.org/stable/90007281

Saunders, A., & Norton, D. A. (2001). Ecological restoration at Mainland Islands in New Zealand. *Biological Conservation, 99*(1), 109-119. doi: https://doi.org/10.1016/S0006-3207(00)00192-0

Scofield, R., Cullen, R., & Wang, M. (2011). Are predator-proof fences the answer to New Zealand's terrestrial faunal biodiversity crisis? *New Zealand Journal of Ecology, 35*(3), 312-317. Retrieved August 12, 2020, from www.jstor.org/stable/24060745

Seabrook-Davison, M. N. H., & Anderson, M. G. (2013). Observation of food presentation behaviour between individual shining cuckoos (Chrysococcyx lucidus) in New Zealand. *Notornis, 60.*

Smith, D. H. V., Wilson, D. J., Moller, H., & Murphy, E. C. (2008). Using artificial nests to explore predation by introduced predators inhabiting alpine areas in New Zealand. *New Zealand Journal of Zoology*, *35*(2), 119-128. https://doi.org/10.1080/03014220809510109

Starling-Windhof, A., Massaro, M., & Briskie, J. V. (2011). Differential effects of exotic predator-control on nest success of native and introduced birds in New Zealand. *Biological Invasions*, *13*(4), 1021-1028. https://doi.org/10.1007/s10530-010-9886-5

Styx Living Laboratory Trust [SLLT]. (n.d.). Styx Mill Conservation Reserve. https://www.thestyx.org.nz/styx-mill-conservation-reserve

Styx Living Laboratory Trust [SLLT]. (2020). *New Zealand eBird; Styx Mill Walkway*. https://ebird.org/newzealand/hotspot/L2484336

Tiritiri Matangi Open Sanctuary. (n.d.). Retrieved from http://www.tiritirimatangi.org.nz/shiningcuckoo

Troup, C. (2015, May 05). Shining cuckoo. Retrieved from https://teara.govt.nz/en/natural-sounds/12499/shining-cuckoo

Wang, Y., Zhou, X., & Engel, B. (2018). Water environment carrying capacity in Bosten Lake basin. *Journal of Cleaner ProductETQq0 0 0 rg BT /Overprint 1 Tf 0 0 1 108.05 263w0(s)-8(e)-5(r)-8(v)23(a)-7(t)-3(i)-7(on)*