EROSION SOIL LOSS AND RECOVERY ON EASTERN NORTH ISLAND HILLCOUNTRY – IMPLICATIONS FOR NUTRIENT MANAGEMENT AND PASTURE PRODUCTIVITY

Brenda Rosser [1,2] and Craig Ross [1]

[1]Landcare Research New Zealand Limited, Private Bag 11052, Palmerston North, New Zealand
[2] Current address: GNS Science, PO Box 30 368, Lower Hutt, New Zealand, b.rosser@gns.cri.nz

Abstract
Landslide scars previously dated (1977, 1961, and 1941) and sampled by Lambert et al. (1984) for pasture production and topsoil characteristics were re-sampled. Pasture dry matter production and selected soil properties were re-measured on the same scars and uneroded control sites for two years, from 2007 to 2009. Results show that after a further 25 years of recovery, no significant increase in pasture production had occurred on the 1941 and 1961 slip scars. Average dry matter pasture production on eroded sites increased from 63% to 78% of pasture production levels on uneroded sites, but improvement was restricted to the youngest 1977 slip scars, where dry matter production increased from 20% to 80% of uneroded levels. Maximum pasture recovery occurred within about 20 years of landsliding and further recovery beyond 80% of uneroded level was unlikely.

The recovery of pasture production on slip scars follows similar recovery to soil physical (e.g. soil depth, particle density, bulk densiy,-N).

on uneroded sites,
e were unable to
variability in total
native forest and
is unlikely total C
of Lambert et al.
oduction levels on
culture on steeper
effects of

c- (r)i7ld lo0wer applicatiy of nutrients
of nutrients to water bodies.

dry matter; carbon; soil recovery; soil
noff

INTRODUCTION

Information on production losses due to soil erosion are essential for quantifying the

Table 1. Chronology of sampling at Te Whanga

Date	Reported by	Measurements
1981	Trustrum & Stephens 1981	Landslip scars mapped and their ages determined
1979–82	Lambert et al 1984	Pasture productivity monitored at 40 sites 7 sites sampled for full soil laboratory characterisation Analysed for total N, organic C, organic P, inorgccP <</MCID 22(o)-2


scars
Total C (%)
Scar age (yrs)
0
1
2
3
4
5
6
7
8
0
20
40
60
80
100
120
140
160

Furthermore, a wider scar age range was studied in the earlier trial, with slip ages ranging from 3 to 74 years, whereas the age range of slips in the latter trial was 27–67 years.

Recovery of topsoil characteristics

The recovery of pasture production was linked to the recovery of topsoil physical and chemical properties of eroded soils by Lambert et al. (1984), who concluded that reduced pasture production on erosion scars was most likely due to the dense, clay-rich substrates, with high bulk de

Trustrum NA, Stephens PR 1981. Selection of hill-country pasture measurement sites by interpretation of sequential aerial photographs. New Zealand Journal of Experimental Agriculture 9: 31–34.

Trustrum NA, Lambert MG, Thomas VJ 1983. The impact of soil slip erosion on hill country pasture production in New Zealand. Proceedings of the Second International Conference on Soil Erosion and Conservation, Honolulu, Hawaii, January 1983.

Trustrum NA, DeRose RC 1988. Soil depth: age relationship of landslides on deforested hillslopes, Taranaki, New Zealand. Geomorphology 1: 143–160.

Trustrum NA, Blaschke PM, DeRose RC, West AW 1990. Regolith changes and pastoral productivity declines following deforestation in steeplands of North Island, New Zealand. Transactions of 14th international Soil Science Congress, Kyoto, Japan, August 1990, Section I: 125–130.

Vincent KW, Milne JDG 1990. Soil pattern or Te Whanga trail site, Wairarapa: soil landscapes, effects of erosion, and soil amelioration in steep eroded siltstone hill country. Unpublished DSIR Land Resources report.

Yadav V, Malanson GP 2009. Modeling impacts of erosion and deposition on soil organic carbon in the Big Creek Basin of southern Illinois. Geomorphology 106: 304–314.

Acknowledgements

We would like to thank Greg Crombie, Manager of Te Whanga Station, for access and helpful discussions about farm management, and Guy Forrester for statistical analysis, Ron DeRose, Mike Marden and Allan Hewitt for reviewing the manuscript and providing useful recommendations, Des Costall for help locating the scars in 2007, and Mike Marden for helping set up the trial. This research was conducted with funding from the Sustainable Land Use Research Initiative (SLURI), FRST contract no. C02X0405.